

”Lifestyle” and Cancer Rates in Former East and West Germany: the possible Contribution of Diagnostic Radiation Exposures

I. Schmitz-Feuerhake and S. Pflugbeil
German Society for Radiation Protection, Berlin, Germany*

***) ingesf@uni-bremen.de**

Introduction

A continuous increase of cancer incidence has been observed in the highly developed countries of Europe and in the U.S. since the second world war. Epidemiologists explain it by “life style” (obesity and unknown factors).

The increase correlates somehow with the rising exposures by X-ray diagnostics and nuclear medicine. Current risk estimates about this influence are usually leading to radiation-induced numbers which are statistically not observable.

In contradiction to the official evaluation there are, however, numerous results of case-control studies in the literature which show measurable contributions of diagnostic exposures.

The risk figures recommended by the ICRP may be too low for several reasons:

- 1) Unjustified use of a DDREF = 2 (Dose and Dose Rate Effectiveness Factor)**
- 2) Uncritical use of the Japanese A-bomb survivor data as a reference.**

Some of the cancers which increased remarkably in the Federal Republic of Germany (FRG) remained much lower in East Germany (GDR) between 1960 and 1989 where diagnostic exposures were applied to a lesser extent because of the national health care system. Low level radiation can cause these diseases and the difference of cancer rates gives rise for renewed evaluation of current risk estimates.

Breast cancer in Germany

Breast cancer in women is predominantly a problem in highly developed countries and in populations of high socioeconomic status.

The incidence in Germany rised continuously until the year 2000 and reached a constant level up to now. The mortality is slowly declining since 1993 which can be explained by improved therapy.

From the 50-ies until the German unification the mortality was about 20 % lower in the GDR (Fig.1).

The female breast is most sensitive for cancer induction by ionising radi-ation which was confirmed by findings after diagnostic X-raying (scoliosis etc.), in stewardesses, and in populations affected by the Chernobyl accident.

There has been no complete registration about X-ray procedures in the FRG and also not in the GDR, and definitely not about the exposures.

Breast Cancer Mortality in Germany




Fig.1

The number of investigations in the GDR has been estimated to 1 per year per inhabitant for 1983-1980 – without dental ones. The FRG estimate is 1.1 for 1978 and 1.3 for 1990-1992.

This difference seems to be not very important. But since 1975 there were rising applications of CT in the FRG with no equivalent in the GDR.

Furthermore, the latter used much less nuclear medicine (I-131 thyroid studies with breast exposures).

A relevant difference between exposures must also have been caused by many "grey" mammographies in the FRG with doses of some 10 mGy.

The very low X-ray energies used in mammography must be considered to be 2-6 fold more effective in inducing cancer than the extremely high energetic gamma radiation which exposed the Japanese A-bomb survivors.

Prostate Cancer

Myles and coworkers (UK) presented a case-control study in 2008 which showed a remarkable effect to the prostate by diagnostic X-rays. After barium enema the risk was elevated by a factor of 2.1 and after hip X-rays by 2.2. They estimated that about 20 % of the current prostate cancer incidence in British males < 60 years are caused by X-raying. The doubling dose can be derived to about 20 mSv from their data.

Other low level effects to the prostate were observed in occupationally exposed British nuclear workers, in pilots, and after the Chernobyl accident.

In Germany, prostate cancer has become the most prevalent cancer disease in men. While the mortality in the GDR was rather constant between 1960 and 1980, it rised at the same time by 50 % in the FRG (Fig.2).




Fig.2

**Prostate Cancer
Mortality**

**in West Germany (FRG)
(above) and
East Germany (GDR)
(below)**

Leukemia in Children


has increased in the FRG by about 50 % between 1960 and 2004 while the GDR showed no elevation (Fig.3).

CTs of the head are supposed to have contributed predominantly to this difference because in childhood up to 30 % of the bone marrow is situated in the skull.

Conclusions

There is evidence from the literature that X-ray diagnostics causes significant contributions to current cancer incidences in the developed countries. This is confirmed by the observed differences between FRG and GDR.

Reduction of diagnostic exposures would be a relevant means for prevention of several prominent cancer diseases.



*Fig.3 Age standardised incidence of childhood leukemia in Western and Eastern Germany (without Berlin)
Data from the German Childhood Cancer Registry in Mainz*

References

- Becker, N., Wahrendorf, J.: Krebsatlas der Bundesrepublik Deutschland 1981-1990. 3. Aufl., Springer 1998
- Bertell, R., Schmitz-Feuerhake, I., Ehrle, L.H.: Pediatric CT research elevates public health concerns: low-dose radiation issues are highly politicized. *Int. J. Health Serv.* 37 (2007) 419-439
- BfS Bundesamt für Strahlenschutz (Ed.): Daten über die Röntgendiagnostik in der ehemaligen DDR. Angerstein, W. et al. BfS-ISH-170/95, Neuherberg 1995
- Hoffmann, D.A. et al.: Breast cancer in women with scoliosis exposed to multiple diagnostic X rays. *J. Natl. Cancer Inst.* 81 (1989) 1307-1312
- Kuni, H., Schmitz-Feuerhake, I., Dieckmann, H.: Mammography screening – neglected aspects of radiation risks. (In German) *Gesundheitswesen* 65 (2003) 443-446
- Myles, P. et al.: Diagnostic radiation procedures and risk of prostate cancer. *Brit. J. Cancer* 98 (2008) 1852-1856
- Pukkala, E. et al.: Breast cancer in Belarus and Ukraine after the Chernobyl accident. *Int. J. Cancer* 119 (2006) 651-8
- Robert Koch Institute and Association of Population-based Cancer Registries in Germany: Cancer in Germany 2005/2006. Incidence and Trends, 7th Edition 2010. Federal Health Reporting. www.gekid.de
- Schmitz-Feuerhake, I. et al.: Estimation of x-ray overexposure in a childhood leukaemia cluster by means of chromosome aberration analysis. *Radiat. Prot. Dos.* 98 (2002) 291-297
- Schmitz-Feuerhake, I., Pflugbeil, S., Pflugbeil, C.: Röntgenrisiko: Abschätzung der strahlenbedingten Meningeome und anderer Spätschäden bei Exposition des Schädels. *Gesundheitswesen* 72 (2010) 246-254
- Schmitz-Feuerhake, I.: Prostatakrebs und diagnostisches Röntgen. *Strahlentelex* Nr. 560-561 vom 6.5.2010, 1-3
- Spix, C., Eletr, D., Blettner, M., Kaatsch, P.: Temporal trends in the incidence rate of childhood cancer in Germany 1987-2004. *Int. J. Cancer* 122 (2008) 1859-1867
- Straume, T.: High-energy gamma rays in Hiroshima and Nagasaki: implications for risk and w_R . *Health Physics* 69 (1995) 954-956
- Vosianov, A.F. et al.: Prostatic intraepithelial neoplasia and apoptosis in benign prostatic hyperplasia before and after Chernobyl accident in Ukraine. *Pathol. Oncol. Res.* 5 (1999) 28-31